



EQUINE DOSING GUIDE

Two intramuscular injections. Four days apart. Maximum of 20 mL per injection site.*

| WEIGHT (LB) | DOSE (mL) |
|-------------|-----------|
| 100 | 1.5 |
| 200 | 3.0 |
| 300 | 4.5 |
| 400 | 6.0 |
| 500 | 7.5 |

| WEIGHT (LB) | DOSE (mL) |
|-------------|-----------|
| 600 | 9.0 |
| 700 | 10.5 |
| 800 | 12.0 |
| 900 | 13.5 |
| 1000 | 15.0 |

| WEIGHT (LB) | DOSE (mL) |
|-------------|-----------|
| 1100 | 16.5 |
| 1200 | 18.0 |
| 1300 | 19.5 |
| 1400 | 21.0 |
| 1500 | 22.5 |

| WEIGHT (LB) | DOSE (mL) |
|-------------|-----------|
| 1600 | 24.0 |
| 1700 | 25.5 |
| 1800 | 27.0 |
| 1900 | 28.5 |
| 2000 | 30.0 |

*Administered at a dose of 3.0 mg/lb (6.6 mg/kg).



EXCEDE® (*ceftiofur crystalline free acid*) Sterile Suspension is indicated for the treatment of lower respiratory tract infections in horses caused by susceptible strains of *Streptococcus equi ssp. zooepidemicus*. EXCEDE is an oil-based substance that may cause some transient swelling and edema around the injection site due to the nature of the compound. Swelling is more likely to occur with injection volumes of more than 15 mLs. In a safety study, swelling completely resolved within 7 days in the majority of cases. Edema may occur a few days after injection. Larger areas of swelling may take longer to resolve. No cases of necrosis, abscess or drainage were reported in any of the studies. EXCEDE is labeled for IM use only.

Important Safety Information: As with all drugs, the use of EXCEDE Sterile Suspension is contraindicated in animals with known allergy to ceftiofur or to B-lactam (penicillins and cephalosporins) group of antimicrobials. Do not use EXCEDE in horses intended for human consumption. The administration of antimicrobials in horses under conditions of stress may be associated with diarrhea, which may require appropriate veterinary therapy. Though safe in cattle when properly administered, inadvertent intra-arterial injection in the ear is possible and is fatal. EXCEDE has a pre-slaughter withdrawal time of 13 days in cattle. Do not use in calves to be processed for veal. For complete details, refer to the full prescribing information.